Effect of ethanol extract of myrmecodia pendens on TGF- β 1 expression and osteoblast cells after tooth extraction (experimental research on cavia cobaya)

Efa Ismardianita^{1*}, Ellyza Nasrul², Yanwirasti³, Mieke Hemiawati⁴

Abstract

Objective: Myrmecodia pendens is often used as a traditional medicine to treat various diseases. A previous study showed that ethanol extract of sarang semut plant (*hypnophytum formicarum* jack) 4.65 mg was effective in fastening establishment of socket granulation tissue after tooth extraction; likewise, the myrmecodia pendens at 3% also fastens the healing process of soft-tissue wound after tooth extraction. The aim of this study is to assess the effect of ethanol extract of 10% myrmecodia pendens on the expression of TGF- β 1 and determine the number of osteoblasts that remain after tooth extraction.

Material and Methods: This study used 24 male marmots divided into 2 groups. All of the groups were extracted in the 1st left-incisor mandibula and 0.1 ml of 0.5% CMC was applied into the socket control group. The treatment group was given 0.1 ml ethanol extract

of myrmecodia pendens 10% dissolved with CMC 0.5%. On day 3, 7, 14 and 21, marmots were decapitated and histopathologic preparations were observed to study the expression of TGF- β 1 and the number of osteoblast cells. Data were analyzed using Kruskal–Wallis test with a probability $p < 0.05$.

Results: Based on the Kruskal–Wallis test, there was a significant difference in the TGF- β 1 expression and the number of osteoblast cells after the administration of ethanol extract of myrmecodia pendens between the control group and treatment group treatment ($p < 0.05$).

Conclusion: The ethanol extract of myrmecodia pendens can fasten the healing of the wound resulting from tooth extraction; it accelerates the healing process by increasing the expression of TGF- β 1 and the number of osteoblast cells.

Keywords: Myrmecodia pendens, Osteoblasts, TGF- β 1, Tooth extraction

Cite this Article: Ismardianita E, Nasrul E, Yanwirasti, Hemiawati M. 2017. Effect of ethanol extract of myrmecodia pendens on TGF- β 1 expression and osteoblast cells after tooth extraction (experimental research on cavia cobaya). *Journal of Dentomaxillofacial Science* 2(3): 150-154. DOI: 10.15562/jdmfs.v2i3.608

¹Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, University Baiturrahmah, Padang, Indonesia

²Department of Clinical Pathologic, Faculty of Medicine, Andalas University, Padang, Indonesia

³Department of Biomedic, Faculty of Medicine, Andalas University, Padang, Indonesia

⁴Departement of Microbiology, Padjajaran University, Bandung, Indonesia

*Corresponding to: Efa Ismardianita, Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, University Baiturrahmah, Padang, Indonesia
efa_ismardianita@yahoo.co.id

Received: 27 August 2017

Revised: 1 October 2017

Accepted: 27 October 2017

Available online: 1 December 2017

Introduction

Tooth extraction often causes wounds. The wound can heal normally, but sometimes it causes some complications that slow down the healing. The rate of likelihood of complications arising from wounds caused by tooth extraction is between 1% and 11.5%¹; research shows that as many as 11 million patients experienced discomfort and complications after tooth extraction, such as pain, swelling, hematoma and difficulties in mastication and speech.² To prevent such complications, dentists made some efforts in finding out the right technique to minimize post-surgical trauma and prescribe medicines appropriately, to treat both systematic or local complications. Antiseptic is the choice for local application in treatment. Antiseptic has several advantages; it kills bacteria and prevents bacteremia; however, there are advantages it using anti-septics, including the difficulty in finding out the right kind of medicinal plant that provides the best possible remedies. Myrmecodia pendens is a medicinal plant. Based on phytochemical test, we found that myrmecodia pendens contains flavonoids,

triterpenoid, phenols and tanin that can be used for anti-inflammation, antibacterial and antioxidant effects.³

The wound healing process that takes place after tooth extraction involves chemotaxis of various cells, cytokines and growing factor. The growing factor responses to early injury on soft and hard tissues include TGF- β 1, BMPs, VEGF, IGFs and PDGF. Transforming growth factor Beta-1 during the inflammation phase initiates and controls chemotaxis, activation and survival of inflammation cells and induces monocytes transformation into macrophages.⁴ Proliferation phase mediates migration and proliferation of endothelial cells, keratinocytes and fibroblasts composed of granulation tissue stimulates VEGF and bFGF angiogenic growth factor expression. Osteogenesis regulates the establishment of bone by inhibiting fibroblasts; it stimulates migration, proliferation, and differentiates mesenchyme cells, transforming them into osteoblasts and increases synthesis and mineralization of bone matrix.^{5,6}

Proliferation of osteoprogenitor cells occurs on day 4, the differentiation of osteoblasts occurs on day 5 and counting is done until day 14. The final stage of osteoblast differentiation and maturation begins on day 15 and lasts until day 28.⁷ In the early stages of osteoprogenitor cell differentiation expression, an increase in TGF- β 1 was detected; otherwise in the final stages of differentiation, the expression of TGF- β decreased.⁸

Material and methods

The extract of ethanol contained the following: Myrmecodia pendens 70%, CMC 0.5%, Ketamin 10% (Ketalar®, Parke Davis) and Xylazine (Bayer), buffer formalin 10%, plank-Rychlo's solution, alcohol, ethanol, xylol, nitrat acid, PBS, xylol, H₂O₂, kit TGF- β 1 (Santa Cruz Biotechnology), biotinylated anti-goat antibody (Dako, Glostrup, Denmark), streptavidin-horseradish peroxidase conjugate, diaminobenzidine (DAB; Sigma Chemical Co.), and HE.

Myrmecodia pendens was brought from Ayawasi village Sorong district, Papua. It was extracted in Chemical Laboratory Research Center Padjajaran University Bandung. Roots of myrmecodia pendens were peeled and sliced to a length of 3-5 mm and were dried (temp 50°C) and grilled with blender. Then a wet mixture containing 250 grams of powder and ethanol 70% (1:5) was prepared. Maceration was performed (shaking the mixture for the first 6 hours and allowing the mixture to settle for the next 18 hours) repeatedly until the filtrate changed color; the filtrate was evaporated using a rotary evaporator (50°C), then fed into the oven until all the water evaporated and the extract became totally dry.

This research is a pure experimental research using post-test control-group design, using 24 cavia cobaya, aged 6-7 weeks. The animals were anesthetized by intramuscular administration of 80 mg/kg Ketamin and 160 mg/kg of Xylazine (1:1). The extraction of the lower incisors was left with needle holder, control group socket was administered 0.1 cc CMC 0.5% and the treatment group was given 0.1 ml of Myrmecodia pendens extract which was dissolved with CMC 0.5%. On days 3, 7, 14 and 21, 3 animals were decapitated and given histopathologic preparations to conduct immune histochemical test, for determining of the levels of TGF- β 1 and HE and osteoblasts; the bodies were observed under light microscope with 40x magnification.

The presence of immunopositive cells according to intensity of staining score was measured based on the following: (I): 0 = no expression;

1 = weak expression; 2 = medium expression; and 3 = strong expression. The presence of immunopositive cells based on distribution was scored as follows: (p): 1 = brown color $\leq$ 10%; 2 = brown >10–50%; and 3 = brown >50–100%. The final score of TGF- β 1 immunopositive cells was calculated by multiplying (p) score and (I) score with the following classification: 0 = no expression; 1: 1–2 = weak positive expression; 2: 3–5 = medium positive expression; and 3: >6 = strong positive expression. The number of osteoblasts was scored as follows: 0 = <10: very less; 1 = 11–50: less; 2 = 51–80: medium; 3 = >80: strong.⁹ Osteoblasts were scored as follows: <10: very less; 11–50: less; 51–80: moderate; and >81: fair.

Results

TGF- β 1 expression was detected as brown color on the cytoplasm, osteoblast membrane cells and a new matrix that was formed (collagen and osteoid). The mean of TGF- β 1 expression of control group and treatment group is presented in figure 1.

TGF- β 1 immunohistochemical expression in control group and treatment group is shown in figure 2.

Based on the results of statistical analysis (Kruskal–Wallis test), we could see there were significant differences in TGF- β 1 expression between the control group and the treatment group; $p = 0.0125$. It can be concluded that the extract ethanol Myrmecodia pendens can increase the expression of TGF- β 1.

The mean number of osteoblasts in control group and treatment group is presented in figure 3.

The number of osteoblast cells in the control group and the treatment group is presented in figure 4.

Results of Kruskal–Wallis test showed there is a significant difference in the number of osteoblasts between the control group and the treatment group;

Figure 1 The mean of TGF- β 1 expression in control group and treatment group

Figure 2 Immunohistochemical view for positive identification of expression of TGF- β 1 on day 3 in collagen mesh (arrow). A. Control group (medium positive), B. Treatment group (strong positive). Day 7 in fibrous tissue, C. control group (strong positive) D. Treatment group (strong positive). Day 14 in osteoblasts. E. Control group (weak positive), F. Treatment group (medium positive). Day 21 in trabecular bone, G. Control group (weak positive), H. Treatment group (weak positive)

Figure 3 The mean of osteoblast cell number in control group and treatment group

p value = 0.0034. Thus, results prove that the extract ethanol myrmecodia pendens can increase the number of osteoblast cells.

Discussion

TGF- β 1 expression in both groups shows a different pattern. On day 3, the TGF- β 1 expression in the treatment group increased and reached its peak. The present study shows results which vary from the previous studies. Expression of TGF- β 1 increased post-trauma and lasted until day 4.^{10,11} The highest level of TGF- β 1 expression occurred

Figure 4 The osteoblast histology observation (arrow) A. Day 3 control group (very less), B. Treatment group (very less), C. Day 7 control group (less), D. Treatment group (less), E. Day 14 control group (good), F. Treatment group (good), G. Day 21 control (good), H. Treatment group (good)

day 3 in the treatment group, which could have been caused by myrmecodia pendens' active substances. Phytochemical test result showed that myrmecodia pendens contained flavonoid, saponin, triterpenoid, tannin and phenolics, all of which could have aided the healing process through different mechanisms.

As an anti-inflammatory agent, flavonoid decreases the permeability of capillary and substracted oedem.¹² Inhibitory functions of arachidonate acid through lipoxygenase and cyclooxygenase pathway cause the inhibition of prostaglandin and leukotriene biosynthesis; as a result, production of prostaglandin, prostacyclin, thromboxane and leukotriene is inhibited so that inflammation is contained.¹⁰ Inflammation phase becomes shorter and comes to the proliferation phase rapidly.¹² Flavonoid inhibits neutrophils degranulation so that it subtracts the extrication of arachidonate acid. Flavonoid could also inhibit leukocytes accumulation in inflammation area, decrease leukocytes adhesion into endothel and reduce the inflammation response.^{13,14}

As antimicrobial agents flavonoid and tannins disturb nucleic acid synthesis and ATPase enzyme activities inhibit metabolism stripe so that bacteria cannot cause further infection and if infection could not be contained the patient's condition will become lethal.^{15,16} Triterpenoid inhibits the growth of bacteria by disturbing the process of formation of bacteria cell membrane.¹⁶ Tannins inhibits transcriptase reverse enzyme and topoisomerase DNA, so that bacteria cannot be formed.¹⁵ Saponin prevents infection by preventing bacteria from lengthening the pro-inflammation cycle.^{17,18}

In the control group, expression increased on day 3, the peak was observed on day 7, on day 14

and day 21 the expression decreased (following a normal healing pattern). The TGF- β 1 distribution and intensity increased after trauma and lasted until day 4, due to the absence of macrophages migration, endothelial cells, keratinocytes, fibroblasts and osteoclasts on the wound area in which the cell secreted TGF- β 1; as the result, TGF- β 1 increased.^{8,9}

TGF- β 1 expression on day 7 in the treatment group is different from Lalani's research¹⁰; on day 4 until day 7 TGF- β 1 expression decreased; meanwhile, in this research we observed that TGF- β 1 expression was still at its peak at this time. The highest level of TGF- β 1 expression observed day 7 shows that giving *Myrmecodia pendens* extract indeed helps fasten the inflammation phase and pushes the inflammation process into proliferation phase. Flavonoid can manage cell function by stimulating the production of TGF- β 1, fibroblasts, keratinocytes, endothelial cells and osteoblasts. This moment is the early stage of osteoblast differentiation.^{5,6,12} During this process of differentiation, TGF- β 1 expression reaches its peak,⁸ which in turn leads to a considerable increase in the proliferation of osteoblasts.¹²

On day 14, TGF- β 1 expression decreases; the said decrease is caused not only by a decrease in the number of chemotactic and mitogenic endotel cells, macrophages, fibroblasts and keratinocytes, but also by various processes such as angiogenesis, epithelization, and fibroplasia; fibroplasia is the last step, that is, it is stage of suppression of inflammation. In the stage of suppression, angiogenesis, epithelization and collagen fibers reach normal levels, and thus proliferation is no longer needed. Some fibroblasts turn to be fibrocytes and others differentiate into osteoblasts.¹²

On day 21, in both groups, TGF- β 1 expression decreased; it could have been caused by the last step of differentiation. In the last step of differentiation and osteoblast maturation (from day 15 to day 28) TGF- β expression begins to decrease because in the last step of differentiation, TGF- β 1 inhibits the proliferation of osteoblasts; as result proliferation of osteoblasts is also reduced.^{8,12,19,20}

Bone tissue regeneration is formed equally between bone resorption by osteoclasts and synthesis by osteoblasts. It correlates with various growth factors. Transforming growth factor induces osteoblast activities, modulating the bone growth.⁴

In this research, the number of osteoblasts in both groups on day 3, 7 and 14 showed the same pattern. The mean number of osteoblasts for the treatment group is higher than that of the control group. On day 21, the number of osteoblasts observed in the treatment group is stable, whereas the number in the control group was still increasing.

On day 3, the mean number of osteoblasts in both groups was very low; this agrees well with the result of osteoprogenitor cell proliferation that happens on day 4; thus, it indicated that the osteoblasts that were found were from the rest of the old bone. On day 4, the number of osteoblasts increased; the mean number of treatment group was higher than that of control group, higher but still low in terms of research criteria. On day 14, the number of osteoblasts increased in both groups. The rate of increase in the number of osteoblasts was very high in the treatment group. It was caused by active substances of *myrmecodia pendens*. Flavonoid can manage cell function by stimulating TGF- β 1 production, whereas TGF- β induces migration and proliferation of osteoblasts.¹³ Triterpenoid can stimulate osteoblast cells to form bone and flavonoid can cause osteoblast differentiation. Accordingly on day 14, the number of osteoblasts was increasing rapidly.

On day 21, in the treatment group, the average number was stable, but in the control group the number kept increasing. The increase might have been caused by TGF- β which still continued to induce osteogenesis process because osteoblast proliferation is still needed to fill the bone defect caused by the wound resulting from tooth extraction.

Conclusion

The application of the ethanol extract *myrmecodia pendens* can assist the wound healing process after tooth extraction by increasing the expression of TGF- β 1. The increase in TGF- β 1 stimulates migration, proliferation, and differentiation of mesenchyme cells into osteoblasts, leading to an increase in the number of osteoblast cells.

Conflict of Interest

The authors report no conflict of interest.

References

1. Simon E, Matee M. Post-extraction complications seen at a referral dental clinic in Dar Es Salaam. *N Dent J* 2001; 51: 273-276.
2. Friedman JW. The prophylactic extraction of third molars: a public health hazard. *Am J Public Health* 2007; 97: 1554-1559.
3. Kareem JJ. Post operative complications associated with non-surgical tooth extraction. *MDJ* 2008; 5: 104-113.
4. Zhao GQ. Consequences of knocking out BMP signaling in the mouse. *Genesis* 2003;35: 43-56.
5. Kasagi S, Chen W. Review TGF-beta1 on Osteoimmunology and the bone component cells. *Cell & Bioscience* 2013; 1-7.
6. Maeda S, Dean DD, Gomez R, et al. The first stage of transforming growth factor beta1 activation is release of

- the large latent complex from the extracellular matrix of growth plate chondrocytes by matrix vesicle stromelysin-1 (MMP-3). *Calcif Tissue Int* 2002;70: 54-65.
7. Dimitriou R, Tsiridis E, Giannoudis PV. Current concepts of molecular aspects of bone healing. *Injury* 2005;36: 1392-1404.
 8. Lalani ZW, Brey EM, Mikos AG, et al. Spatial and temporal localization of transforming growth factor-beta 1, bone morphogenetic protein-2 and platelet-derived growth factor-A in healing tooth extraction sockets in rabbit model. *J Oral Maxillofac Surg* 2003;61: 1061-1072.
 9. Mohamed MAH, Younis WH, Yaseen NY. The effect of autologous bone marrow-derived stem cells with estimation of molecular events on tooth socket healing in diabetic rabbits (immunohistochemical study). *J Bagh College Dentistry* 2013;25.
 10. Sabir. Pemanfaatan flavonoid di bidang kedokteran gigi. *Majalah kedokteran Gigi Edisi Khusus Temu Ilmiah Nasional III* 2003: 81-83.
 11. Nijveldt RJ, Nood E, Hoorn DEC, et al. Flavonoids: a review of probable mechanism of action and potential application. *Am J Clin Nutr* 2001;74: 418-425.
 12. Lakhanpal P, Rai DK. Quercetin: a versatile flavonoid. *IJMU* 2007;2: 22-37.
 13. Akiyama H, Kazuyasu F, Osamu Y, et al. Antibacterial action of several tannins against staphylococcus aureus. *J Antimicrob Chemot* 2001;48: 487-491.
 14. Ajizah, A. Sensitifitas salmonella typhimurium terhadap ekstrak daun psidium guajava l. *Journal Bioscient* 2004; 1: 31-38.
 15. Gautam GK, Mishra SH, Kumar A, et al. Phytochemical evaluation of the ethanol extract of baubinia tomentosa Linn. (Root). *IJPLS* 2012;3: 1675-1676.
 16. Robinson T. "The basic of higher plants 6th ed". In Disadur PK. *Kandungan organik tumbuhan tinggi*. Bandung: ITB; 1995.
 17. Huang Z, Nelson ER, Smith RL, et al. The sequential expression profiles of growth factors. *J Tissue Eng* 2007;13: 2311-2320.
 18. Iwata J, Hosokawa R, Sanchez-Lara PA, et al. Transforming growth factor-beta regulates basal transcriptional regulatory machinery to control cell proliferation and differentiation in cranial neural crest-derived osteoprogenitor cells. *J Biol Chem* 2010;285: 4975-4982.
 19. Spinella-Jaegle S, Roman-Roman S, Faucheu C, et al. Opposite effects of bone morphogenetic protein-2 and transforming growth factor-beta1 on osteoblast differentiation. *Bone* 2001;29: 323.
 20. Zhang Z, Nakashima K, Zhou X, et al. The novel zinc finger-containing transcription factor osterix is required for osteoblast differentiation and bone formation. *Cell* 2008;108: 17-29.

This work is licensed under a Creative Commons Attribution